

Alerts, Notices, and Case Reports

Peritoneal Coccidioidomycosis Diagnosed Incidentally at Herniorrhaphy

JOSE A. PEREZ, Jr, MD
EDWARD L. ARSURA, MD
Bakersfield, California

INFECTION WITH THE pathogenic fungus, *Coccidioides immitis*, is a frequent event in areas endemic to this organism. Most infections are asymptomatic or remain localized to the lungs, and in only about 0.5% of host encounters with *C immitis* does the organism produce extrapulmonary symptoms.¹ In the Central Valley of California in the most recent 12-month period, there has been an outbreak of *C immitis* infections without historical parallel.² One uncommon presentation of *C immitis* infection is coccidioidal peritonitis. This disorder can present with diffuse abdominal pain or as an abdominal wall hernia.^{3,4}

Report of a Case

The patient, a 30-year-old man, noticed periumbilical swelling three months before admission. Two months before admission, he detected a right inguinal swelling. He otherwise felt well without fever or constitutional symptoms. A chest x-ray film showed no abnormalities.

During an operation, an umbilical hernia was repaired without incident. During right inguinal herniorrhaphy, however, the hernia sac was found to be thickened and edematous. Straw-colored fluid was noted in the hernia sac, and this revealed a leukocyte count of 4.2×10^9 per liter (4,200 per μ l), with 0.80 (80%) lymphocytes, 0.07 (7%) neutrophils, 0.01 (1%) basophils, 0.07 (7%) monocytes, and 0.01 (1%) mesothelial cells. Examination of the inguinal hernia sac revealed granulomata with *C immitis* spherules visible on Gomori-methenamine silver stain. Serologic studies for *Coccidioides immitis* revealed a complement-fixing antibody titer of 1:256. The patient did well with a course of intravenous amphotericin B colloidal dispersion.

Discussion

Coccidioidal peritonitis was first described in 1896 by Roxford and Gilchrist in an autopsy report,⁵ and since that time, 13 cases of coccidioidal peritonitis have been reported. Curiously, as in our patient, 3 of 7 patients in the series of Saw and co-workers had no evidence of

pulmonary disease, and 2 of these presented with right inguinal hernia.³ In a review by Chen, 4 of 13 patients presented with uncomplicated inguinal hernia.⁵

Consideration to increasingly unusual presentations of coccidioidal infection should be given as the number of cases reaches epidemic proportions.

REFERENCES

1. Stevens D: *Coccidioides immitis*, In Mandell GL, Douglas RG Jr, Bennett JE (Eds): Principles and Practice of Infectious Disease, 3rd edition. New York, NY, Churchill Livingstone, 1990, pp 2008-2017
2. Centers for Disease Control and Prevention: Coccidioidomycosis—United States, 1991-1992. MMWR 1993; 42:21-24
3. Saw EC, Shields SJ, Comer TP, Huntington RN: Granulomatous peritonitis due to *Coccidioides immitis*. Arch Surg 1974; 108:369-371
4. Weisman IM, Moreno AJ, Parker AL, Sippo WC, Liles WJ: Gastrointestinal dissemination of coccidioidomycosis. Am J Gastroenterol 1986; 81:589-593
5. Chen KTK: Coccidioidal peritonitis. Am J Clin Pathol 1983; 80:514-516

Acute Renal Failure Following Intravenous Streptokinase Infusion for Acute Myocardial Infarction

YOCHAI BIRNBAUM, MD
BORIS STRASBERG, MD
ELDAD RECHAVIA, MD
YORAM NEUMAN, MD
BRACHA STAHL, Mpharm
Petah-Tikva, Israel

STREPTOKINASE, ADMINISTERED EITHER intravenously or by the intracoronary route, is one of the drugs most frequently used for thrombolysis in acute myocardial infarction. Acute renal failure is a relatively rare complication following thrombolytic therapy with streptokinase.¹ We describe the case of a patient in whom acute renal failure developed nine days after thrombolytic therapy with streptokinase for acute inferior wall myocardial infarction.

Report of a Case

The patient, a 62-year-old man, was admitted to hospital because of acute inferior wall myocardial infarction. In the past he had had hypercholesterolemia and a diaphragmatic hernia. On physical examination he was in good general condition, his blood pressure was 130/80 mm of mercury, and his heart rate was 82 beats per minute. The lungs were clear, and there were no signs of

(Birnbbaum Y, Strasberg B, Rechavia E, Neuman Y, Stahl B: Acute renal failure following intravenous streptokinase infusion for acute myocardial infarction. West J Med 1993 Apr; 158:406-409)

(Perez JA Jr, Arsura EL: Peritoneal coccidioidomycosis diagnosed incidentally at herniorrhaphy. West J Med 1993 Apr; 158:406)

From the Department of Medicine, Kern Medical Center, Bakersfield, California, University of California, Los Angeles, School of Medicine.

Reprint requests to Edward L. Arsura, MD, Chair, Department of Medicine, Kern Medical Center, 1830 Flower St, Bakersfield, CA 93305.

From the Cardiac Intensive Care Unit (Dr Birnbbaum), Departments of Cardiology (Drs Birnbbaum, Strasberg, and Rechavia) and Internal Medicine A (Dr Neuman), and the Drug Information Center (Dr Stahl), Beilinson Medical Center, Petah-Tikva, and the Sackler School of Medicine, Tel-Aviv University, Tel-Aviv, Israel.

Reprint requests to Yochai Birnbbaum, MD, Cardiac Intensive Care Unit, Beilinson Medical Center, Petah-Tikva, Israel 49 100.